

June 1992

LBL-32324

UCB-PTH-92/15

ANALYSIS OF RUNNING COUPLING CONSTANT UNIFICATION IN STRING THEORY ¹

Mary K. Gaillard and Rulin Xiu

Department of Physics, University of California

and

Physics Division, Lawrence Berkeley Laboratory, 1 Cyclotron Road

Berkeley, CA 94720

Abstract

We use recently obtained 2-loop string coupling constants to analyze a class of string models based on orbifold compactification. Assuming weak coupling at the string scale m_s and single-scale unification leads to restrictions on the spectrum of massive ($M_Z \ll M \leq m_s$) matter supermultiplets and/or on the Kac-Moody algebra level.

¹This work was supported by the Director, Office of Energy Research, Office of High Energy and Nuclear Physics, Division of High Energy Physics of the U.S. Department of Energy under contract DE-AC03-76SF00098 and in part by the National Science Foundation under grant PHY-90-21139.

In the past few years, progress in string phenomenology has been very encouraging. On one hand, the elucidation of the effective theories that describe the low energy dynamics of the light string excitations has been carried out to one genus and higher genus [1],[2] in string calculations. A systematic scheme has been developed to obtain an effective theory with modular invariance to all loop orders. One loop and all loop gauge couplings in string effective theories have been obtained for a class of models [3]–[7]. On the other hand, the construction of “realistic” string models has been developed to produce some explicit string models. Even though none of them are totally realistic, some are very close to the standard model and might even make some good predictions [8]. Because of this progress, it is now possible confront specific models with data.

In this paper, we discuss the running gauge coupling constant unification in string effective theories. It is interesting to note that string unification has, in general, a more flexible meaning than standard model unification. On the other hand, because of the relation between the unification scale and the gauge coupling constant in string effective theory, the string analysis is more restrictive. We restrict our discussion to the minimal string model from an $N = 1$ orbifold with no $N = 2$ space-time supersymmetry. In this case, the running coupling constants do not have nontrivial moduli-dependent string threshold corrections [2]. By minimal string model, we mean that the model has $SU(3) \times SU(2) \times U(1)$ as observable gauge group (that is, we neglect possible mixing with additional $U(1)$ ’s, and do not consider intermediate scale gauge symmetry breaking) and $N=1$ supersymmetry below the compactification scale. In addition to the standard model matter content there are generally extra particles which become massive at some high energy scales. Our analysis will impose restrictions on the spectra of massive states. We pay particular attention to this kind of model not only because it is simple to analyse but also because it is the closest approximation to the standard model to date.

In the following we give the two-loop renormalization group equations (RGE) in string effective theory for a class of models derived from orbifold compactification. We take as a starting point the result of [3] that suggests that two-loop unification occurs at the string scale m_s . We will see that, under the assumption of weak coupling at

the string scale, the scale $\tilde{m}_P = \frac{1}{2}m_P e^{\frac{1}{2}}$, where m_P is the Planck mass, can be taken as the effective string unification scale in the first order RGE.

Whereas the unification of the running gauge coupling constants at some scale is a coincidence in the standard model, a relation among gravitational and gauge coupling constants at the string scale is required in string theory. It has been shown [9] that at the string scale $m_s^2/m_P^2 = \alpha_i k_i$, $m_s = 2/\sqrt{\alpha'}$. Here the Planck mass m_P is defined by $G_N = \kappa^2/8\pi = 1/m_P^2$. α' is the string tension and k_i is the Kac-Moody level of the gauge algebra G_i . For a nonabelian gauge group its level is an integer and is determined by the corresponding Kac-Moody algebra realized on a world sheet. For a $U(1)$ gauge group the level is defined to be $Y = k_1/2$ [10], where Y is the double-pole coefficient in the operator product expansion of the corresponding Kac-Moody currents. The relation is obtained under the assumption that any gauge symmetry has its origin in an internal Kac-Moody algebra as is the case for all known closed string theories. It is a general result in the sense that it does not depend on the particular realization of a gauge algebra in a model. To see the dependence of coupling constants on a scale, a string loop calculation has to be applied. In the following analysis, the $U(1)$ charge is the weak hypercharge of the standard model, but normalized as in $SU(5)$, so that $k_1 = k_2 = k_3$ corresponds to coupling constant unification in the usual sense.

In string effective theory, the gauge coupling constant g is determined by the vacuum expectation value of the dilaton s : $g^{-2} = \langle Res \rangle$. The one loop modular invariant gauge coupling has been calculated [3] for orbifolds with gauge group $E_8 \otimes E_6 \otimes U(1)^2$ and $k_a = 1$. A correspondence between the effective coupling constant and the running coupling constant in quantum field theory was proposed in [3], where it was shown that the relation can be interpreted as the two loop running coupling constant [11] with the string scale m_s as the two loop unification scale, up to possible string threshold corrections. More precisely, the results of [3] suggest the identification $g_a^2(\frac{1}{2}g^2 M_P^2) = g^2/k_a - \epsilon_a/4\pi$, where $M_P = m_P/\sqrt{8\pi}$ is the reduced Planck mass, and ϵ_a is a small threshold correction. Here the running coupling constant $g^2(\lambda^2)$ is to be interpreted as the measured coupling constant with external momenta $-p^2 = \lambda^2$. This is related [12] to the scale μ of the $\overline{MS}$ regularization scheme by

$\lambda^2 = e\mu^2$. In this scheme the unification scale becomes $\mu_s^2 = e^{-1}\alpha m_P^2/4 = \alpha\tilde{m}_P^2$. Consequently, for scales $\mu > M_{SUSY}$, the running gauge coupling constants in string phenomenology can be written as:

$$g_a^{-2}(\mu) = g_a^{-2} - \frac{C_a^G}{8\pi^2} \ln 2 - \frac{1}{8\pi^2} (3C_a^G - C_a^M) \ln \frac{\mu_s}{\mu} + \frac{C_a^G}{8\pi^2} \ln \frac{g_a^2}{g_a^2(\mu)} + \frac{1}{8\pi^2} \sum_A C_a^A \ln \frac{Z_A(\mu_s)}{Z_A(\mu)} - \frac{1}{2} \sum_I \beta_a^I \ln [\eta^2(it^I) \overline{\eta^2(it^I)} (t + \bar{t})^I]. \quad (1)$$

Here t^I are the moduli and $4e\mu_s^2/m_P^2 = \alpha_a k_a = \alpha$. The index A denotes the matter content: $C_a^M = \sum_A C_a^G$. C_a^G is the quadratic Casimir operator in the adjoint representation of the gauge subgroup labeled by a , and $C_a^A = \text{Tr}(T_a^A)^2$, where T_a^A is any generator of the gauge group G_a for the representation R_A , is related to the quadratic Casimir operator $C_2^a(R_A)$ by $\dim(G_a)C_a^A = \dim(R_A)C_2^a(R_A)$. The matter wave-function normalization satisfies at one loop

$$\frac{d \ln Z_A(\mu)}{d \ln \mu} = \sum_b C_2^b(R_A) \frac{\alpha_b}{\pi}. \quad (2)$$

Now we write eq.(1) as:

$$\alpha_a^{-1}(\mu) = k_a \alpha^{-1} + \frac{b_a}{2\pi} \ln \frac{\tilde{m}_P}{\mu} + \delta_a^{(2)} + \Delta_a, \quad b_a = C_a^M - 3C_a^G. \quad (3)$$

Here $\delta_a^{(2)}$ is the “second order” correction; integration of (2) using the one-loop expression for $\alpha(\mu)$ gives, for scales $\mu > M_{SUSY}$:

$$\delta_a^{(2)} = \frac{b_a}{2\pi} \ln \alpha - \epsilon_a + \frac{C_a^G}{2\pi} \ln \frac{\alpha_a}{\alpha_a(\mu)} + \sum_{A,b} \frac{1}{\pi b_b} C_a^A C_2^b(R_A) \ln \frac{\alpha_b(\mu_s)}{\alpha_b(\mu)}, \quad (4)$$

where $\epsilon_a = C_a^G \ln 2/2\pi$, and Δ_a is the string threshold correction [1]:

$$\Delta_a = -2\pi \sum_I \beta_a^I \ln [\eta^2(it^I) \overline{\eta^2(it^I)} (t + \bar{t})^I]. \quad (5)$$

Eq.(3) expresses the result that in first order the running coupling constants approximately “unify ” at Planck scale with some string threshold corrections.

Although the result (1) was explicitly derived [3] for the $E_8 \otimes E_6 \otimes U(1)^2$ orbifold, its generalization to other Z_N orbifolds with $k_a = 1$ is straightforward. We will also

consider the assumption that it can be extended to $k_a \neq 1$. Here we restrict our analysis to orbifolds with no $N = 2$ supersymmetry sectors; in this case [2]

$$\beta_a^I = 0 = \Delta_a. \quad (6)$$

RGE analyses including moduli-dependent threshold corrections have been considered elsewhere [13]. The restriction (6) implies a condition [1]–[3] on the spectrum of gauge nonsinglets and their modular weights.

For the minimal string model, the above results then give.

$$\alpha_a^{-1}(M_Z) = k_a \alpha^{-1} + \frac{b_a^{(I)}}{2\pi} \ln \frac{M_{SUSY}}{M_Z} + \frac{b_a^{(II)}}{2\pi} \ln \frac{\tilde{m}_P}{M_{SUSY}} + \Pi_a + \delta_a^{(2)}. \quad (7)$$

Here

$$\Pi_a = \frac{1}{2\pi} \sum_E b_{aE} \left(1 + \frac{\alpha_a}{2\pi} C_a^G + \sum_b \frac{\alpha_b}{\pi} C_a^E C_b(R_E) \right) \ln \frac{\mu_s}{m_E}, \quad (8)$$

where we have used the first order expression for $\ln[\alpha_a(\mu_s)/\alpha(m_E)]$ in the second order terms. The index E extends over extra, massive particles.

$$b_a^{(I)} = \begin{pmatrix} 4.2 \\ -3 \\ -7 \end{pmatrix}$$

is the β -function after supersymmetry is broken (we assume two Higgs doublets with mass $m_H \approx M_Z$), and

$$b_a^{(II)} = \begin{pmatrix} 6.6 \\ 1 \\ -3 \end{pmatrix}$$

is the β -function in the supersymmetric region for the minimal supersymmetric standard model (MSSM).

Note that in string phenomenology, the parameters α , M_{SUSY} , $\ln(\mu_s/m_E)$, M_W are supposed to be determined dynamically. Then the coupling constants at the weak scale can in principle be computed. In addition, according to the most popular supersymmetry breaking mechanism, gaugino condensation [14], M_{SUSY} is determined

by the gauge group in the hidden sector. Then eq.(7) implies that low energy measurements can in principle put constraints on both the hidden sector gauge group and the observable gauge group. Unfortunately, there is not as yet a satisfactory theory of supersymmetry breaking by hidden gaugino condensation; we will use the phenomenological constraint that M_{SUSY} must be within about an order of magnitude of a TeV , and define $M_T = 1TeV$ as the “first order” SUSY breaking scale.

We first consider the case $k_a = 1$; presently most string models produced are of this kind. We note that the result (3) was obtained [3] from the analysis of the one-loop chiral anomaly, which is connected by supersymmetry to the conformal anomaly, in the context of supergravity. In the context of supersymmetric Yang-Mills theory, the authors of Ref. [11] argue that the two-loop condition (3) should be valid to all loop orders; their argument is related to the well-known result that the chiral anomaly is determined completely at the one loop level. We therefore minimize the approximations in the evaluation of (3); specifically, we do not use the lowest order expression for $\ln[\alpha(\mu_s)/\alpha(M_{SUSY})]$. In addition to the one-loop approximation used in writing the last term in (4), we make the following approximations. We use the lowest order expression for $\ln[\alpha(M_{SUSY})/\alpha(M_W)]$ to evaluate $\alpha(M_{SUSY})$. This gives, using LEP data [15] and other results [16]:

$$\alpha^{-1}(M_{SUSY}) + b_a^{(I)} z = \tilde{B}_a, \quad z = \frac{1}{2\pi} \ln \frac{M_{SUSY}}{M_T}, \quad (9)$$

where the quantities

$$\tilde{B} = \begin{pmatrix} 57.25 \pm .11 \\ 30.84 \pm .11 \\ 11.23 \pm .80 \end{pmatrix}, \quad b_a^{(I)} = \begin{pmatrix} 4.30 \\ -2.86 \\ -7.23 \end{pmatrix},$$

include the two-loop corrections as obtained in this approximation. We used as input the values quoted in [16], but with a larger error ($\delta\alpha_3^{-1} = \pm.79$) on α_s which covers the slightly higher recent result [17]. We then evaluate (3) taking $\mu = M_{SUSY}$, and approximating $\ln \alpha_a(M_{SUSY}) = -\ln \tilde{B}_a + \tilde{B}_a^{-1} b_a^{(I)} z$, which is justified by the assumption $.1 < M_{SUSY}/M_T < 10$, $|z| \leq .37$, and $\ln \alpha_a(\mu_s) = \ln \alpha$, which is justified (for $k_a = 1$) under our weak coupling assumption: $\alpha^{-1} \gg \epsilon_a = (0, .22, .33)$. This

gives:

$$\alpha^{-1} + \frac{b_a^{(I)} - b_a^{(II)}}{2\pi} \ln \frac{M_{SUSY}}{M_T} + \Pi_a = B_a + c_a \ln \alpha, \quad (10)$$

$$B = \begin{pmatrix} 18.87 \pm .11 + .03\eta \\ 19.98 \pm .11 + .04\eta \\ 27.01 + .82\eta \end{pmatrix}, \quad c_a = \begin{pmatrix} 1.11 \\ 1.53 \\ -.106 \end{pmatrix},$$

where $\eta = \pm 1$ is the sign of $\delta\alpha_3^{-1}$.

We are interested in finding constraints on string phenomenology from low energy measurements. We write (9) in the form:

$$\alpha^{-1} - 1.11 \ln \alpha - 2.57z + \Pi_1 = B_1, \quad (11)$$

$$\alpha^{-1} - 1.53 \ln \alpha - 4.09z + \Pi_2 = B_2, \quad (12)$$

$$\alpha^{-1} + .106 \ln \alpha - 4.40z + \Pi_3 = B_3. \quad (13)$$

These equations have no solution for $\Pi_a = 0$. In this case the analysis reduces to that of MSSM which has been done [18] to second order and gives: $M_{GUT} = 10^{16 \pm 0.3} \text{GeV}$ and $\alpha_{GUT} \simeq 26$. In string coupling constant unification, we have an additional constraint: $\alpha_{GUT} m_P = M_{GUT} = \mu_s$, which cannot be satisfied.

However, the universal anomaly cancellation condition (6) generally requires the existence of additional particles, so we expect $\Pi_a \neq 0$. Since $\Pi_a \geq 0$, if we impose the constraint $|z| \leq .37$, each of eqs. (11)–(13) gives an upper bound on α^{-1} ; the strongest bound is from (11):

$$\alpha^{-1} \leq 16.83 \quad (14)$$

(Note that although the first term on the right hand side of (4) is not strictly speaking of second order, this bound, together with the weak field assumption $\alpha \leq 1$ assures that it, as well as the threshold correction ϵ_a , is comparable to the second order corrections.)

Next consider the difference

$$\Pi_3 - \Pi_2 = 7.03 - 1.64 \ln \alpha + .31z \pm .11 + .78\eta \geq 6.03 \quad (15)$$

using the weak coupling constraint $\alpha \leq 1$. This tells us that for this kind of model to work, at least one extra $SU(3)$ triplet is required. Note that (15) is independent of k_1 , so our analysis excludes the 3-generation $SU(3) \times SU(2) \times U(1)$ model of [8], which is a level one model with no extra $SU(3)$ triplets. For general orbifold compactifications based on the heterotic string, the chiral multiplets in the observable sector are in $27 + \overline{27}$ of E_6 . The massive states come in particle-anti-particle pairs that we denote according to their $(SU(3)_c \times SU(2)_L)_{|Q|}$ quantum numbers (Q = electric charge) as

$$q = (3, 2), \quad \ell = (1, 2), \quad d = (3, 1)_{\frac{1}{3}}, \quad u = (3, 1)_{\frac{2}{3}}, \quad e = (1, 1)_1.$$

Then

$$\Pi_a = \sum_i b_a^i \Pi_i, \quad \Pi_i = \sum_{M_i} \frac{n_i}{\pi} \ln \frac{\mu_s}{M_i} \geq 0,$$

where n_i is the number of pairs with mass M_i . Using the first order expressions for the β -function constants b_a^i , Eq.(8), gives

$$\begin{aligned} \Pi_1 &= .1\Pi_q + .8\Pi_u + .2\Pi_d + .3\Pi_\ell + .6\Pi_e, \\ \Pi_2 &= 1.5\Pi_q + .5\Pi_\ell, \quad \Pi_3 = \Pi_q + .5\Pi_u + .5\Pi_d. \end{aligned} \tag{16}$$

Writing

$$\begin{aligned} \Pi_1 - .4\Pi_3 + .2\Pi_2 &= 12.06 - .8\alpha^{-1} + 1.46 \ln \alpha + 1.63z \pm .13 - .29\eta \\ &= .6\Pi_u + .4\Pi_\ell + .6\Pi_e \geq 0, \end{aligned} \tag{17}$$

gives a stronger bound on α :

$$\alpha^{-1} \leq 11.84. \tag{18}$$

Using the weak coupling constraint, we can also exclude models in which the heavy states all transform as $(3, 1) + (\bar{3}, 1) + (1, 2)$:

$$\begin{aligned} \Pi_1 - .4\Pi_3 - .6\Pi_2 &= -1.2\Pi_q + .6\Pi_u + .6\Pi_e \\ &= -3.92 + .23 \ln \alpha - 1.64z \pm .18 - .32\eta \leq -2.81, \quad \Pi_q \geq 2.34, \end{aligned} \tag{19}$$

for $\alpha \leq 1$. In addition we get, using (18),

$$\begin{aligned} \Pi_3 - \frac{2}{3}\Pi_2 &= .5 \left(\Pi_u + \Pi_d - \frac{2}{3}\Pi_\ell \right) \\ &= 13.69 - \frac{1}{3}\alpha^{-1} - 1.13 \ln \alpha \pm .07 + .79\eta + 1.67z \quad \begin{cases} \geq 11.06 \\ \leq 15.42 \end{cases}. \end{aligned} \quad (20)$$

which in turn implies that $(3, 2) + (\bar{3}, 2)$ states do not make the full contribution to Π_3 . Using (18) and $\alpha \leq 1$, together with the above results, gives

$$\begin{aligned} \Pi_2 &\leq 20.54, \quad \Pi_3 \leq 28.41, \\ 2.34 &\leq \Pi_q \leq 13.69, \quad 22.12 \leq \Pi_d + \Pi_u \leq 52.14. \end{aligned} \quad (21)$$

The above bounds have implications for the masses of these states; assuming that Π_q is dominated by the lightest multiplet, we find, for example

$$1.3GeV \leq m_q \leq 3.9 \times 10^{15}GeV. \quad (22)$$

Clearly the lower mass range is ruled out; if the upper bound on Π_q is saturated, we require $n_q \geq 2$, which in turn implies $m_q \geq 2.8 \times 10^9 GeV$. The value found for $\Pi_d + \Pi_u$ has to be interpreted in terms of a lower bound on the number of these states: $n_u + n_d \geq 2$. If the upper bound on $\Pi_d + \Pi_u$ is saturated, we require $n_u + n_d \geq 5$; the corresponding mass bounds are

$$m_{u \text{ or } d} \leq 5TeV, \text{ if } n = 2, \quad m_{u \text{ or } d} \geq 36TeV \text{ if } n \geq 5. \quad (23)$$

Finally, we consider models with arbitrary levels: $k_a \geq 1$. Although no minimal string model with a higher Kac-Moody level has as yet been constructed, the basic technique has already been established [10], and there seems to be no fundamental difficulty in obtaining such a model. If the low energy effective theory differs from the $k_a = 1$ case only by the additional factors k in the couplings of the gauge supermultiplets to the dilaton s , then eq.(3) should still be valid. There is an additional threshold-like effect from the last term in (4): $\ln \alpha_b(\mu_s) = \ln \alpha - \ln k_b$.

If the model is clever enough, it may be possible to achieve universal anomaly cancellation without extra particles. We will only consider this more restrictive case;

then the low energy measurements constrain the values of k_1, k_2 and k_3 . First we note that for models that unify to a larger gauge group at the string scale, $\alpha_1^{-1} = \alpha_2^{-1} = \alpha_3^{-1} = k\alpha^{-1}$, we would have the constraint: $M_{GUT} = \mu_s = k\alpha_{GUT}m_P \geq \alpha_{GUT}m_P$, which cannot be satisfied without additional particles or nontrivial string threshold corrections or both. For the models we are considering, which do not have nontrivial string threshold corrections, extra $SU(3)$ triplets are again required for a model to work.

Setting $\Pi_a = 0$ eqs.(11)–(13) now appear as:

$$k_1\alpha^{-1} - 1.11 \ln \alpha - 2.56z = B_1(k), \quad (24)$$

$$k_2\alpha^{-1} - 1.53 \ln \alpha - 4.09z = B_2(k), \quad (25)$$

$$k_3\alpha^{-1} + .106 \ln \alpha - 4.41z = B_3(k), \quad (26)$$

$$B_a(k) = B_a + \sum_{A,b} \frac{1}{\pi b_b} C_a^A C_2^b(R_A) \ln k_b.$$

If $\ln k_a \sim 1$, the additional correction is of same order as the uncertainty in the SUSY threshold. Using the constraints $-.37 \leq \alpha \ln \alpha \leq 0$ for $0 \leq \alpha \leq 1$ and $|z| \leq .37$, (25) and (26) give

$$1.16 \leq \frac{k_3}{k_2} \leq 2.32, \quad (27)$$

if we restrict the integers $1 \leq k_2, k_3 \leq 10$. This allows for the possibility $k_2 = 2, k_3 = 1$; however, the solution to (25) and (26) with these values gives $9.5 \leq \alpha^{-1} \leq 12$, $M_{SUSY} \leq 3GeV$, which is excluded. For fixed $k_3/k_2 = 2$ and $k_2\alpha^{-1} \geq 4.2$, z decreases as k_2 increases, so there is no solution with $k_3/k_2 = 2$; in a similar way one can exclude the cases $k_3/k_2 = 7/4, 5/3, 8/5, \dots$. We find solutions with $k_2 = 5, k_3 = 7, 3 \leq \alpha^{-1} \leq 4.6, 4.1 \leq k_1 \leq 4.7, -1.3 \leq z \leq .8$, and with $k_2 = 6, k_3 = 8, 3 \leq \alpha^{-1} \leq 4.7, 4.8 \leq k_1 \leq 5.3, -.59 \leq z \leq 2$, which cover the allowed range for M_{SUSY} , and are the only solutions with $k_3/k_2 = 7/4$ and $4/3$, respectively. In addition we find that $k_2 = 2, k_3 = 3, \alpha^{-1} \approx 9.3, k_1 \approx 1.8$, and $k_2 = 4, k_3 = 6, \alpha^{-1} \approx 4.5, k_1 \approx 4$, are allowed within the errors but with marginally acceptable values for z : $M_{SUSY} \geq 7.15TeV$ in the first case and $M_{SUSY} \leq 140GeV$ in the second case; these are the only solutions with $k_3/k_2 = 3/2$.

True coupling constant unification with $k_a = 1$ is of course the most attractive possibility. The relation between the coupling and the scale at unification in orbifold compactification is inconsistent with the values determined [18] from a fit of the MSSM to the data. This could be evaded without introducing extra particles in several ways. In Calabi-Yau compactification, it is generally assumed that the unification scale is determined by the vacuum expectation value of a modulus field t rather than the dilaton s , in which case the relation between the coupling constant and the unification scale would not hold. Since these theories have not been studied to the same extent as orbifold models, no conclusion can really be drawn. Another possibility is that the MSSM is valid to the scale of $10^{16} GeV$, above which there is a larger gauge group; in this case nothing can be inferred from the data on the value of $\alpha = m_s^2/m_P^2$. A third alternative is orbifold models with moduli-dependent threshold corrections [13]. However in these models, the moduli-dependence of the effective potential for gaugino condensation [19] breaks the continuous symmetries that would otherwise protect [20] scalar particles from acquiring large masses when local supersymmetry is broken. For this reason, models with universal anomaly cancellation may be more attractive for phenomenology. We have seen that these models can fit the data, which in turn provide restrictions on the spectrum of heavy gauge nonsinglet supermultiplets; in particular, there must be color triplet (3,1) and (3,2) states. Some of these could be light enough to be produced in collider experiments, and, for a suitable mass range, these states could have interesting implications for proton stability and/or baryogenesis. A similar analysis including moduli-dependent threshold corrections and extended gauge groups will be given elsewhere.

Acknowledgement.

This work was supported in part by the Director, Office of Energy Research, Office of High Energy and Nuclear Physics, Division of High Energy Physics of the U.S. Department of Energy under Contract DE-AC03-76SF00098 and in part by the National Science Foundation under grant PHY-90-21139.

References

- [1] L. Dixon, V. Kaplunovsky and J. Louis, *Nucl. Phys.* **B355**: 649 (1991).
- [2] I. Antoniadis, K.S. Narain and T.R. Taylor *Phys. Lett.* **B267**: 37 (1991).
- [3] Mary K. Gaillard, T.R. Taylor, LBL-31464 UCB-PTH-91/65 NUB-3039 (1991) to be published in *Nucl. Phys.* **B**.
- [4] J.P. Derendinger, S. Ferrara, C. Kounnas, and F. Zwirner, *Phys. Lett.* **B271**: 307 (1991).
- [5] J.P. Derendinger, S. Ferrara, C. Kounnas, and F. Zwirner, CERN-TH.6004/91, LPTENS-91-4 (1991).
- [6] G.L. Cardoso and B.A. Ovrut, *Nucl. Phys.* **B369**: 351 (1992).
- [7] P. Binetury, G. Girardi, and R. Grimm, *Phys. Lett.* **B265**: 111 (1991).
- [8] A. Font, L.E. Ibáñez, F. Quevedo, and A. Sierra, *Nucl. Phys.* **B331**: 421 (1990).
- [9] P. Ginsparg, *Phys. Lett.* **B197**: 139 (1987).
- [10] A. Font, L.E. Ibáñez, F. Quevedo, and A. Sierra, *Nucl. Phys.* **B345**: 389 (1990).
- [11] M.A. Shifman and A.I. Vainshtein, *Nucl. Phys.* **B277**: 456-486 (1986).
- [12] see, *e.g.*, T. Muta, *Foundations of Quantum Chromodynamics, An Introduction to Perturbative Methods in Gauge Theories* (World Scientific, Singapore, 1987).
- [13] L.E. Ibáñez, D. Lüst and G.G. Ross, *Phys. Lett.* **B272**: 251 (1991); L.E. Ibáñez and D. Lüst, CERN-TH.6380/92 (1992).
- [14] H.P. Nilles, *Phys. Lett.* **115B**: 193 (1982).
- [15] The LEP Collaborations: ALEPH, DELPHI, L3 and OPAL, *Phys. Lett.* **B276**: 247 (1992);

- [16] P. Langacker, UPR-0492T, Invited talk given at Int. Workshop on Electroweak Physics Beyond the Standard Model, Valencia, Spain, Oct 2-5, 1991.
- [17] The ALEPH collaboration, CERN-PPE/92-33.
- [18] U. Amaldi, W. de Boer, and H. Furstenuau, *Phys. Lett.* **B260**: 447 (1991); P. Langacker and M.-X. Luo, *Phys. Rev. D* **44**: 817 (1991).
- [19] A. Font, L.E. Ibáñez, D. Lüst and F. Quevedo, *Phys. Lett.* **B245**: 401 (1990); S. Ferrara, N. Magnoli, T. Taylor and G. Veneziano, *Phys. Lett.* **B245**: 409 (1990); M. Cvetič, A. Font, L.E. Ibáñez, D. Lüst and F. Quevedo, *Nucl. Phys.* **B361**: 194 (1991); T. Taylor, *Phys. Lett.* **B252**: 59 (1990); P. Binétruy and M.K. Gaillard, *Phys. Lett.* **253B**: 119 (1991); M.K. Gaillard, A. Papadopoulos and D. Pierce, *Phys. Rev.* **D45**: 2057 (1992); D. Lüst and C. Munős, *Phys. Lett.* **B279**: 272 (1992).
- [20] P. Binétruy and M.K. Gaillard, *Phys. Lett.* **232B**: 83 (1989) and *Nucl. Phys.* **B358**: 121 (1991).